

# PATTERNS OF TRAUMATIC BRAIN INJURIES IN PATIENTS PRESENTING AT A TERTIARY CARE UNIT

PIR SAAD, MUHAMMAD SHEHZAD KHAN WAZIR, DAWOOD SOHAIL, SHAHBAZ ALI KHAN,  
ATIF IBRAHIM, MUHAMMAD WAQAS

*Ayub Medical College, Abbottabad*

*Corresponding author Address: Opposite Lady Garden Park, Regional Tax Office, Main Mansehra Road,  
Abbottabad, KPK, Pakistan. Contact no: 0321-9963813*

## ABSTRACT

**Background:** Traumatic brain injury is a change in function of the brain or alternatively any other evidence of brain abnormality that is caused by a sudden external force. The purpose of this study was to evaluate the distribution of traumatic brain injuries at Ayub Teaching Hospital. The results could be used to give recommendations for the betterment of approach towards and management of traumatic brain injuries care.

**Methods:** Questionnaire based data was collected from an inpatient population of patients who presented to the Neurosurgical Unit of Ayub Teaching Hospital with traumatic brain injuries. CAT scan was used as the imaging modality for preliminary diagnosis. The data was collected over a period of three years. Non-probability purposive sampling was used as the sampling technique. Patients of both sexes and all ages were included in the study.

**Results:** Out of a total of 1938 patients, 1470 (75.9%) were males and 468 (24.1%) were females. Patients from 20 to 40 years age group (38.1%) represented the greatest number. Patients from Abbottabad (41.8%) were found to be the most numbered contributors. Fall (52.6%) was found to be the most common external cause of TBIs followed by Road Traffic Accidents (34.1%). Most of the patients (42.4%) had a GCS score falling between 8 and 12. No lesion was found at the initial CAT scan in most of the patients (27.2%). Depressed Skull Fracture (21.4%) was the most common abnormal finding in initial imaging. Conservative (78.3%) treatment was provided to most of the patients keeping in view the appropriate management requirements. 97.8% of the patients were treated successfully.

**Conclusion and Recommendations:** The quality of care at Neurosurgery Ayub Teaching Hospital was found to be up to the mark for traumatic brain injuries patients. However, the standard of care at Kashmir and Balakot needs to be re-evaluated. At the same time meta-analysis outcomes should be established for the spectrum of the different aspects of traumatic brain injuries patients. More studies should be conducted to evaluate the factors responsible for making males more vulnerable to TBIs relative to females.

**Keywords:** Trauma, Glasgow Coma Scale, Computerized Automated Tomography scan, Extradural hematoma, Subdural hematoma, Lesion, Subarachnoid hemorrhage, Intracranial bleed, Linear Skull Fracture, Depressed Skull Fracture

## INTRODUCTION

A traumatic brain injury can be defined as a nondegenerative, noncongenital insult to the brain from an external mechanical force, possibly leading to permanent or temporary impairment of cognitive, physical, and psychosocial functions, with an associated diminished or altered state of consciousness<sup>1</sup>. Traumatic brain injury is a substantial contributor of mortality and morbidity throughout the world<sup>2</sup>. Traumatic brain injuries cost an average of \$56 billion annually in U.S<sup>3</sup> while in Europe, approximately 1.6 million people are admitted to the hospital every year with traumatic brain injuries<sup>4</sup>.

In Pakistan, the number of deaths attributed to traumatic brain injuries is increasing day by day<sup>5</sup>. The annual incidence of traumatic brain injuries in Pakistan has been estimated to be 50/100,000 with bread earning youth contributing the most to this figure<sup>6</sup>. The situation in Pakistan is particularly alarming because with the greater increase in the number of road traffic accidents and violence gaining its hold in the society, trauma patients are increasing in number substantially<sup>7</sup>. A World Bank study established the fact that each year almost four percent of Pakistan's total population is pushed below poverty due to health shocks<sup>8</sup>. Social re-emergence of a traumatic patient, both physically as well

as mentally is a big problem in our setup<sup>9</sup>. The prognostic model of Traumatic Brain Injuries for a developing country like Pakistan differs significantly from the developed world and hence based on epidemiological studies more valid and suitable prognostic and management models should be come up with<sup>10</sup>

The purpose of this study was to bring into light the distribution of traumatic brain injuries and to come up with recommendations to enrich research, ameliorate patient care and improve prevention of trauma. The pattern of traumatic brain injuries was followed in a number of variables like age, gender, area of residence, type of trauma, Glasgow Coma Scale (GCS), preliminary diagnosis, management and outcome. The study was conducted at Ayub Teaching Hospital, Abbottabad, a 1000 bed tertiary care facility.

## MATERIALS AND METHODS

It was a snapshot study that was conducted at the Neurosurgery Unit of Ayub Teaching Hospital. One thousand nine hundred and thirty eight patients were included in the study. Duration of the study was 3 years i.e. from May, 2012 to June, 2015. Non-probability

purposive sampling was used as the sampling technique. Neurosurgical in patients who presented with brain injury secondary to trauma and only those patients who had no abnormality or a single abnormality on imaging out of the predefined lesions were included in the study. Outdoor patients were excluded from the study.

A detailed questionnaire that was filled after taking informed consent from the patients or their attendants was used as the data collection tool. Preliminary diagnosis was made by CAT scan as the imaging modality. Glasgow Coma Scale (GCS) score was used as a mode of estimation of consciousness of the patients and the extent of their brain damage. The data was processed by SPSS-16. Descriptive statistics were calculated. Variables were expressed by calculating frequency and percentage calculation.

## RESULTS

Out of a total of 1938 patients, 624 (32.2%) were less than 13 years of age, 318 (16.4%) between 13 and years of age, 738 (38.1%) 20 to 40 years old while 258(13.3%) patients were more than 40 years old.

**Table-1:** Age of Patients

Age Group (Years)	Frequency	Percent	Valid Percent	Cumulative Percent
Less than 13 years old	624	32.2	32.2	32.2
13 to 19 years old	318	16.4	16.4	48.6
20 to 40 years old	738	38.1	38.1	86.7
More than 40 years old	258	13.3	13.3	100.0
Total	1938	100.0	100.0	

1470(75.9%) were males and 468(24.1%) were found to be females out of the total 1938 patients.

**Table-2:-** Gender

Gender	Frequency	Percent	Valid Percent	Cumulative Percent
Male	1470	75.9	75.9	75.9
Female	468	24.1	24.1	100.0
Total	1938	100.0	100.0	

**Table-3:-** Area of Residence

Area of Residence	Frequency	Percent	Valid Percent	Cumulative Percent
Abbottabad	810	41.8	41.8	41.8
Haripur	132	6.8	6.8	48.6
Mansehra	468	24.1	24.1	72.8
Batagram	228	11.8	11.8	84.5
Northern Areas	48	2.5	2.5	87.0
Kashmir	216	11.1	11.1	98.1
Swat	12	.6	.6	98.8
Afghanistan	12	.6	.6	99.4
Balakot	12	.6	.6	100.0
Total	1938	100.0	100.0	

Greater number of patients presented at night and in evening timings as compared to morning and day timings.

**Table 4:** Time of Admission

Time of admission	Frequency	Percent	Valid Percent	Cumulative Percent
5:00 am to 11:59 am	300	15.5	15.5	15.5
12:00 pm to 5:00 pm	402	20.7	20.7	36.2
5:01 pm to 10:00 pm	678	35.0	35.0	71.2
10:01 pm to 4:59 am	558	28.8	28.8	100.0
Total	1938	100.0	100.0	

**Table 5:** Cause of Trauma

Cause of Trauma	Frequency	Percent	Valid Percent	Cumulative Percent
Fall	1020	52.6	52.6	52.6
Road Traffic Accident	660	34.1	34.1	86.7
Assault	204	10.5	10.5	97.2
Firearm Injury	54	2.8	2.8	100.0
Total	1938	100.0	100.0	

**Table 6:** Glasgow Coma Scale at the time of presentation

Glasgow Coma Scale	Frequency	Percent	Valid Percent	Cumulative Percent
13-15	660	34.1	34.1	34.1
8-12	822	42.4	42.4	76.5
5-7	444	22.9	22.9	99.4
3-4	12	.6	.6	100.0
Total	1938	100.0	100.0	

**Table 7:** Preliminary Diagnosis on CAT Scan

Preliminary Diagnosis on CAT Scan	Frequency	Percent	Valid Percent	Cumulative Percent
No Lesion	528	27.2	27.2	27.2
Extra Dural Hematoma	258	13.3	13.3	40.6
Subdural Hematoma	54	2.8	2.8	43.3
Intracranial Bleed	234	12.1	12.1	55.4
Linear Skull Fracture	294	15.2	15.2	70.6
Depressed Skull Fracture	414	21.4	21.4	92.0
Subarachnoid Hemorrhage	156	8.0	8.0	100.0
Total	1938	100.0	100.0	

**Table 8:** Management

Management	Frequency	Percent	Valid Percent	Cumulative Percent
Conservative	1518	78.3	78.3	78.3
Operated	420	21.7	21.7	100.0
Total	1938	100.0	100.0	

**Table 9:** Outcome

Outcome	Frequency	Percent	Valid Percent	Cumulative Percent
Alive	1896	97.8	97.8	97.8
Dead	42	2.2	2.2	100.0
Total	1938	100.0	100.0	

The vast majority of patients i.e. 810(41.8%) were from District Abbottabad. Patients from Swat, Afghanistan and Balakot contributed the least to the figure with 12(0.6%) from each of these areas. There

were 132(6.8%) patients from District Haripur, 468(24.1%) from District Mansehra, 228(11.8%) from Batagram, 48(2.5%) from Northern areas including Gilgit Baltistan and 216(11.1%) from Kashmir.

Fall was found to be the most frequent contributor to traumatic brain injuries i.e. 1020(52.6%) patients presented with a history of fall. This was followed by Road Traffic Accidents contributing a total of 660(34.1%) patients. 204(10.5%) patients had a history of assault while only 54(2.8%) patients had Firearm injury as the cause of Traumatic Brain Injury.

On Glasgow Coma Scale, 660(34.1%) patients scored from 13 to 15. The most frequent GCS was found to be between 8 and 12 i.e. 822(42.4%) patients fell in this category at the time of their presentation. 444(22.9%) patients had a GCS score of 5 to 7 while 12(0.6%) patients were graded as 3 to 4 on Glasgow Coma Scale at the time of their presentation.

1518(78.3%) of the patients were treated conservatively while 420(21.7%) had to be operated on as part of their management.

Patients with no lesion i.e. 528(27.2%) had the lion's share. This was followed by Depressed Skull Fracture i.e. 414(21.4%) patients out of the total had a depressed skull fracture. 258(13.3%) patients were found to be positive for an Extra Dural Hematoma while only 54(2.8%) patients had a Subdural Hematoma as their diagnosis on CAT Scan. 234(12.1%) patients had an Intracranial Bleed, 294(15.2%) Linear Skull Fracture and 156(8%) patients had Subarachnoid Hemorrhage as their preliminary diagnosis.

The outcome was found to be exceptionally good with 1896(97.8%) being treated successfully while a meager number of 42(2.2%) out of 1938 couldn't make it back to life.

## DISCUSSION

A physical injury to the living tissue caused by an external source is called a trauma<sup>11</sup>. Traumatic brain injury is therefore a physical injury to the brain that is not caused by a progressive process, is acquired and is caused by an external mechanical insult that may cause a long term or short term abnormality of behavioral, physical, social and/or psychological functions with an associated impaired consciousness<sup>12</sup>.

Age was found to be an important factor while taking into consideration the epidemiology of Traumatic Brain Injuries. Majority of the patients were found to be below 40 years of age. This is in consistence with a study performed by John Bruns<sup>13</sup>. As most of the patients were relatively young, the outcome was positive for majority of the patients. This is in accordance with the study by Hilaire J<sup>14</sup>.

As per this study majority of the patients were found to be males. This is in agreement with a study performed in 2004 in which 121 previous studies were critically studied for the prevention, incidence and risks

of traumatic brain injuries<sup>15</sup>. The finding of males being more sufferers of traumatic brain injuries is also strongly supported by the Center for Disease Control and Prevention (CDC), USA, according to which Traumatic Brain Injuries rates are higher in males in every age group as compared to females<sup>16</sup>. A study conducted by Kraus J in 1984 also supports this statistical outcome<sup>17</sup>. Men below 40 years of age are more vulnerable to Traumatic Brain Injuries<sup>18</sup>. Males are in majority when it comes to contribution to the number of total TBI patients<sup>19</sup>.

Regarding the different areas from which TBI patients presented it can be presumably established that the greatest influx of traumatic brain injuries patients was from the nearest locations. This is probably due to the logical fact that patients were being treated at their respective nearest primary and secondary care units decreasing the number of patients from far flung areas at our unit. Despite of the weak practical translation of healthcare hierarchy in Pakistan<sup>20</sup>, it can be assumed that our tertiary care unit had a much lesser load of TBI patients because majority of the patients were being treated at their respective areas' local healthcare facilities. However, TBI patients from Kashmir and Balakot surpassed the rest of the areas in number taking into account the relative distances and the time that it takes to reach our healthcare unit from the different areas studied.

As far as time of the day that has the greatest load of TBI patients is concerned, the outcome is dispersed. However evening and night timings are definitely ahead of morning timings in numbers as evident by the data.

Fall was found to be the most frequent external source of injury in our setup followed by Road Traffic Accidents and Assault. Surprisingly our region shared the trend with USA as the same distribution was found by a survey conducted from 2002-2006 at USA by the Center for Disease Control and Prevention<sup>21</sup>. Fall was also found to be the most frequent contributor to Traumatic Brain Injuries by a survey conducted at Tennessee and the number was found to be on the rise from 1996-2010<sup>22</sup>. However, by comparison and contrast of different studies, it can be concluded that the external source of TBI varies depending on the geography and a fixed or nearly consistent pattern cannot be established between two different geographical regions or even within the region.

Glasgow Coma Scale (GCS) for majority of the patients was found to be more than 8 and that had an impact on the positive outcome of majority of the patients. This is consistent with a study performed by Cisu D in 1997<sup>23</sup>.

Linear Skull Fracture was found to be the most frequent finding on CAT Scans of patients presented with TBIs while Subdural Hematoma was found to be the least contributor. The use of CAT scan in establishing appropriate diagnosis and planning management accordingly is of utmost importance<sup>24</sup>.

Majority of the patients were treated conservatively in view of their GCS findings at the time of their presentation and diagnosis. This was in agreement with a study conducted in 2008 by GH Yattoo who concluded that only 10% to 20% of TBI patients need surgical management while the rest benefit from nonsurgical management options<sup>25</sup>.

With majority of the patients falling in a category of younger age i.e. below 40 years<sup>26 27</sup>, GCS score of more than 8<sup>28</sup> and most of the patients requiring conservative treatment, the outcome for a tremendous number of patients was good as expected.

## CONCLUSION

Majority of TBI patients are below 40 years of age. Males are more vulnerable to TBIs as compared to females. The greatest numbers of patients receive treatment from their nearest healthcare facilities. Fall by far is the most common external cause of TBI followed by road traffic accidents. GCS correlates positively with the outcome and most of the patients fall between a GCS score of 8 and 12. Most of the patients have no lesion at the time of their preliminary diagnosis on CAT scan while depressed skull fracture is the most common abnormality of all on CAT scans at the time of their presentation. Majority of the patients are managed conservatively and the results are substantially positive for a great number of patients. With the exceptionally great number of patients being treated successfully at Ayub Teaching Hospital, it can be safely concluded that the quality of care for traumatic brain injuries patients is up to the mark at the Neurosurgery Unit of ATH.

## RECOMMENDATIONS

Specific trauma centers with staff specifically trained for traumatic brain injuries should be established in Northern parts of the country including Abbottabad. The healthcare units at Kashmir and Balakot should be looked into for flaws and thus should be corrected accordingly. Data regarding traumatic brain injuries is lacking in Pakistan. Statistical analysis of traumatic brain injuries should be conducted at regular intervals of time and therefore standardized requirements should be met according to the needs of different localities. More studies should be conducted and thus an accurate and reliable prognostic scoring system should be introduced

keeping in view the patterns of traumatic brain injuries identified.

## REFERENCES

1. Emedicine.medscape.com. Traumatic Brain Injury (TBI) - Definition and Pathophysiology: Overview, Epidemiology, Primary Injury [Internet]. 2016 [cited 9 January 2016]. Available from: <http://emedicine.medscape.com/article/326510-overview>
2. Heegaard W, Biros M. Traumatic Brain Injury. *Emergency Medicine Clinics of North America*. 2007;25(3):655-678.
3. Ninds.nih.gov. Traumatic Brain Injury: Hope Through Research [Internet]. 2016 [cited 9 January 2016]. Available from: [http://www.ninds.nih.gov/disorders/tbi/detail\\_tbi.htm](http://www.ninds.nih.gov/disorders/tbi/detail_tbi.htm)
4. Exchange C. Brain Injury: Long term outcome after traumatic brain injury | International Encyclopedia of Rehabilitation [Internet]. Cirrie.buffalo.edu. 2016 [cited 9 January 2016]. Available from: <http://cirrie.buffalo.edu/encyclopedia/en/article/338/>
5. Medforum.pk. Head Injury and its Consequences – a One Year Study in Karachi [Internet]. 2016 [cited 9 January 2016]. Available from: <http://www.medforum.pk/index.php/article-database/9-articles/44-head-injury-and-its-consequences-a-one-year-study-in-karachi>
6. Umerani MS e. Traumatic brain injuries: experience from a tertiary care centre in Pakistan. - PubMed - NCBI [Internet]. Ncbi.nlm.nih.gov. 2016 [cited 9 January 2016]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24535786>
7. Jpma.org.pk. Trauma Care in Pakistan [Internet]. 2016 [cited 9 January 2016]. Available from: [http://www.jpma.org.pk/full\\_article\\_text.php?article\\_id=1337](http://www.jpma.org.pk/full_article_text.php?article_id=1337)
8. Hafeez M. Poverty and Poor Health in Pakistan: Exploring the Effects of Privatizing Healthcare - Harvard International Review [Internet]. Harvard International Review. 2014 [cited 9 January 2016]. Available from: <http://hir.harvard.edu/poverty-and-poor-health-in-pakistan/>
9. Jpma.org.pk. The decrepit state of trauma care in under-developed healthcare settings and the need to integrate trauma rehabilitation as a continuum of care [Internet]. 2016 [cited 9 January 2016]. Available from:

- http://jpma.org.pk/full\_article\_text.php?article\_id=7150
9. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. BMJ. 2008;336(7641):425-429.
10. Merriam-webster.com. Definition of TRAUMA [Internet]. 2016 [cited 9 January 2016]. Available from: <http://www.merriam-webster.com/dictionary/trauma>
11. Emedicine.medscape.com. Traumatic Brain Injury (TBI) - Definition and Pathophysiology: Overview, Epidemiology, Primary Injury [Internet]. 2016 [cited 9 January 2016]. Available from: <http://emedicine.medscape.com/article/326510-overview>
12. [Internet]. 2016 [cited 9 January 2016]. Available from: <http://onlinelibrary.wiley.com/doi/10.1046/j.1528-1157.44.s10.3.x/pdf>
13. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2367127/>
14. Incidence, Risk Factors And Prevention Of Mild Traumatic Brain Injury: Results Of The Who Collaborating Centre Task Force On Mild Traumatic Brain Injury. J REHABIL MED. 2004;(43):28-60.
15. [Internet]. 2016 [cited 9 January 2016]. Available from: [http://www.cdc.gov/traumaticbraininjury/pdf/BlueBook\\_factsheet-a.pdf](http://www.cdc.gov/traumaticbraininjury/pdf/BlueBook_factsheet-a.pdf)
16. Kraus J, Black M, Hessol N, LEY P, Rokaw W, Sullivan C et al. The Incidence Of Acute Brain Injury And Serious Impairment In A Defined Population. American Journal of Epidemiology [Internet]. 1984 [cited 9 January 2016];119(2):186-201. Available from: <http://aje.oxfordjournals.org/content/119/2/186.short>
17. About.com Men's Health. Why Men Suffer More Acquired Brain Injury [Internet]. 2016 [cited 9 January 2016]. Available from: <http://menshealth.about.com/library/bltrauma.htm>
18. Aans.org. The American Association of Neurological Surgeons [Internet]. 2016 [cited 10 January 2016]. Available from: <http://www.aans.org/patient%20information/conditions%20and%20treatments/traumatic%20brain%20injury.aspx>
19. Waheed M. WHO EMRO | Health system strengthening | Programmes | Pakistan [Internet]. Emro.who.int. 2016 [cited 9 January 2016]. Available from: <http://www.emro.who.int/pak/programmes/health-system-strengthening-hss.html>
20. [Internet]. 2016 [cited 9 January 2016]. Available from: [http://www.cdc.gov/traumaticbraininjury/pdf/BlueBook\\_factsheet-a.pdf](http://www.cdc.gov/traumaticbraininjury/pdf/BlueBook_factsheet-a.pdf)
21. [Internet]. 2016 [cited 9 January 2016]. Available from: [https://tn.gov/assets/entities/health/attachments/ECodes\\_1996-2010.pdf](https://tn.gov/assets/entities/health/attachments/ECodes_1996-2010.pdf)
22. Cifu D, Keyser-Marcus L, Lopez E, Wehman P, Kreutzer J, Englander J et al. Acute predictors of successful return to work 1 year after traumatic brain injury: A multicenter analysis. Archives of Physical Medicine and Rehabilitation. 1997;78(2):125-131.
23. Stiell I, Wells G, Vandemheen K, Clement C, Lesiuk H, Laupacis A et al. The Canadian CT Head Rule for patients with minor head injury. The Lancet. 2001;357(9266):1391-1396.
24. Yattoo G, Tabish A. The profile of head injuries and traumatic brain injury deaths in Kashmir. J Trauma Manage Outcomes. 2008;2(1):5.
25. Abiebr.com. Key Points | ERABI [Internet]. 2016 [cited 10 January 2016]. Available from: <http://www.abiebr.com/key-points-0>
26. Braininjury.com. brain injury .com | Traumatic Brain Injury In The Aging Population [Internet]. 2016 [cited 10 January 2016]. Available from: <http://www.braininjury.com/tbi-and-the-elderly.shtml>
27. Sevil Bilgin B. Gait and Glasgow Coma Scale scores can predict functional recovery in patients with traumatic brain injury. Neural Regeneration Research [Internet]. 2012 [cited 10 January 2016];7(25):1978. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4298893/>